arXiv:0910.0506v1 [math.DG] 3 Oct 2009

Bifurcations of Wavefronts on an r-corner II: Semi-local classification

Takaharu Tsukada [1]

College of Humanities & Sciences, Department of Mathematics,
Nihon University

Abstract

We give a classification of generic bifurcations of intersections of wavefronts generated by different points of a hypersurface with or without boundaries.

1 Introduction

In [11] we investigated the theory of *reticular Legendrian unfoldings* which describes bifurcations of wavefronts on a hypersurface germ with a boundary, a corner, or an r-corner in a smooth manifold and studied the stabilities and a generic classification of such bifurcations.

In this paper we give a semi-local classification of bifurcations of wavefronts generated by a hypersurface with an r-corner. The wavefronts generated around several points on the initial hypersurface intersect. For example we consider the below figures in which the initial wavefront with a boundary in a plane and wavefronts generated by the initial wavefront and the boundary to normal directions are described respectively. The generated wavefronts bifurcate and typical shapes of bifurcations occur. The ${}^1({}^0A_1{}^0A_2)$, ${}^1({}^0A_1{}^0A_1)$, and ${}^1({}^0A_1{}^0B_2)$ fronts are intersections of fronts generated by different points of the initial wavefront. The purpose of this paper is the classification of such generic bifurcations of intersections of wavefronts.



Figure 1: The initial wavefront V^0 with a boundary and generated wavefronts ($e = \emptyset$, $t_1 < t_2 < t_3$)

As the bifurcation theory of wavefronts are described by reticular Legendrian unfoldings, we shall introduce the notion of *multi-reticular Legendrian unfoldings* in order to describe the theory of generic bifurcations of intersections of wavefronts generated by different points of a hypersurface with an r-corner.

---

[1] Higashijujo 3-1-16 Kita-ku, Tokyo 114-0001 JAPAN. e-mail : tsukada@math.chs.nihon-u.ac.jp

The main result of this paper is a classification of generic bifurcations of intersections of wavefronts generated by a hypersurface without boundary or with boundaries in a manifold with dimension 2 and 3. The classification list of generating families of generic multi-reticular Legendrian unfoldings are given before Theorem 5.6. We also draw all figures of such bifurcations at the last of this paper.

This paper consists of four sections. In section 2 we investigate the theory of map germs with respect to *the reticular* t-$(\mathcal{P}$-$\mathcal{K})_{(m)}$*-equivalence relation* which plays important roles as generating families of multi-reticular Legendrian unfoldings. In section 3, we review the theory of reticular Legendrian unfoldings which is developed in [11]. In section 4 we introduce the notion of multi-reticular Legendrian unfoldings and investigate their equivalence relation, generating families, and stabilities. In section 5 we reduce our investigation to finitely dimensional jet spaces and give a generic classification of multi-reticular Legendrian unfoldings.

2 Stabilities of unfoldings

In this section we investigate the theory of map germs with respect to *the reticular* t-$(\mathcal{P}$-$\mathcal{K})_{(m)}$*-equivalence relation* which is proved by almost parallel methods of [10].

We denote by $\mathcal{E}(r;k_1,r;k_2)$ the set of all germs at 0 of smooth maps $\mathbb{H}^r\times\mathbb{R}^{k_1}\to\mathbb{H}^r\times\mathbb{R}^{k_2}$ and set $\mathfrak{M}(r;k_1,r;k_2)=\{f\in\mathcal{E}(r;k_1,r;k_2)|f(0)=0\}$. We denote $\mathcal{E}(r;k_1,k_2)$ for $\mathcal{E}(r;k_1,0;k_2)$ and denote $\mathfrak{M}(r;k_1,k_2)$ for $\mathfrak{M}(r;k_1,0;k_2)$.

If $k_2=1$ we write simply $\mathcal{E}(r;k)$ for $\mathcal{E}(r;k,1)$ and $\mathfrak{M}(r;k)$ for $\mathfrak{M}(r;k,1)$. Then $\mathcal{E}(r;k)$ is an $\mathbb{R}$-algebra in the usual way and $\mathfrak{M}(r;k)$ is its unique maximal ideal. We also denote by $\mathcal{E}(k)$ for $\mathcal{E}(0;k)$ and $\mathfrak{M}(k)$ for $\mathfrak{M}(0;k)$. We remark that $\mathcal{E}(r;k,p)$ is an $\mathcal{E}(r;k)$-module generated by p-elements..

We denote by $J^l(r+k,p)$ the set of l-jets at 0 of germs in $\mathcal{E}(r;k,p)$. There are natural projections:

$$\pi_l:\mathcal{E}(r;k,p)\longrightarrow J^l(r+k,p), \pi^{l_1}_{l_2}:J^{l_1}(r+k,p)\longrightarrow J^{l_2}(r+k,p)\ (l_1>l_2).$$

We write $j^lf(0)$ for $\pi_l(f)$ for each $f\in\mathcal{E}(r;k,p)$.

Let $(x,y)=(x_1,\cdots,x_r,y_1,\cdots,y_k)$ be a fixed coordinate system of $(\mathbb{H}^r\times\mathbb{R}^k,0)$. We denote by $\mathcal{B}(r;k)$ the group of diffeomorphism germs $(\mathbb{H}^r\times\mathbb{R}^k,0)\to(\mathbb{H}^r\times\mathbb{R}^k,0)$ of the form:

$$\phi(x,y)=(x_1\phi^1_1(x,y),\cdots,x_r\phi^r_1(x,y),\phi^1_2(x,y),\cdots,\phi^k_2(x,y)).$$

We denote by $\mathcal{B}_n(r;k+n)$ the group of diffeomorphism germs $(\mathbb{H}^r\times\mathbb{R}^{k+n},0)\to(\mathbb{H}^r\times\mathbb{R}^{k+n},0)$ of the form:

$$\begin{aligned}\phi(x,y,u)=(&x_1\phi^1_1(x,y,u),\cdots,x_r\phi^r_1(x,y,u),\\ &\phi^1_2(x,y,u),\cdots,\phi^k_2(x,y,u),\phi^1_3(u),\ldots,\phi^n_3(u)).\end{aligned}$$

We denote $\phi(x,y,u)=(x\phi_1(x,y,u),\phi_2(x,y,u),\phi_3(u))$, $\frac{\partial f_0}{\partial y}=(\frac{\partial f_0}{\partial y_1},\cdots,\frac{\partial f_0}{\partial y_k})$, and denote other notations analogously.

Lemma 2.1 (cf. [13, Corollary 1.8]) *Let B be a submodule of $\mathcal{E}(r;k+n+m',m)$, A_1 be a finitely generated $\mathcal{E}(m')$-submodule of $\mathcal{E}(r;k+n+m',m)$ generated d-elements, and A_2 be a* *finitely generated $\mathcal{E}(n+m')$ submodule of $\mathcal{E}(r;k+n+m',m)$. Suppose*

$$\mathcal{E}(r;k+n+m',m) = B + A_2 + A_1 + \mathfrak{M}(m')\mathcal{E}(r;k+n+m',m) \\ +\mathfrak{M}(n+m')^{d+1}\mathcal{E}(r;k+n+m',m).$$

Then

$$\mathcal{E}(r;k+n+m',m) = B + A_2 + A_1,$$
$$\mathfrak{M}(n+m')^{d}\mathcal{E}(r;k+n+m',m) \subset B + A_2 + \mathfrak{M}(m')\mathcal{E}(r;k+n+m',m).$$

We say that $f_0 = (f_{0,1},\ldots,f_{0,m})(x,y)$, $g_0 = (g_{0,1},\ldots,g_{0,m})(x,y) \in \mathcal{E}(r;k+n)$ are *reticular $\mathcal{K}_{(m)}$-equivalent* if there exist $\Phi_i \in \mathcal{B}_n(r;k+n)$ and a unit $\alpha_i \in \mathcal{E}(r;k+n)$ such that $g_{0,i} = \alpha_i \cdot f_{0,i} \circ \Phi_i$ for $i = 1,\ldots,m$. We denote $\Phi = (\Phi_1,\ldots,\Phi_m)$, $\alpha = (\alpha_1,\ldots,\alpha_m)$ and we call (Φ,α) a reticular $\mathcal{K}_{(m)}$-isomorphism from f to g. We remark that f_0 and g_0 are reticular $\mathcal{K}_{(m)}$-equivalent if and only if $f_{0,i}$ and $g_{0,i}$ are reticular $\mathcal{K}$-equivalent for $i = 1,\ldots,m$.

Lemma 2.2 *Let $f_0(x,y) \in \mathfrak{M}(r;k,m)$ and $z = j^l f_0(0)$. Let $O^l_{r\mathcal{K}}(z)$ be the submanifold of $J^l(r+k,m)$ consist of the image by π_l of the orbit under the reticular $\mathcal{K}_{(m)}$-equivalence of f_0. Put $z = j^l f_0(0)$. Then*

$$T_z(O^l_{r\mathcal{K}}(z)) = \pi_l((\langle f_{0,1}, x\frac{\partial f_{0,1}}{\partial x}\rangle_{\mathcal{E}(r;k)} + \mathfrak{M}(r;k)\langle\frac{\partial f_{0,1}}{\partial y}\rangle) \\ \times\cdots\times(\langle f_{0,m}, x\frac{\partial f_{0,m}}{\partial x}\rangle_{\mathcal{E}(r;k)} + \mathfrak{M}(r;k)\langle\frac{\partial f_{0,m}}{\partial y}\rangle)).$$

We say that a map germ $f_0 = (f_{0,1},\ldots,f_{0,m})(x,y) \in \mathfrak{M}(r;k,m)$ is *reticular $\mathcal{K}_{(m)}$-l-determined* if all map germ in $\mathfrak{M}(r;k,m)$ which has the same l-jet of f_0 is reticular $\mathcal{K}_{(m)}$-equivalent to f_0.

Lemma 2.3 (Cf., [10, Lemma 2.3]) *Let $f_0 = (f_{0,1},\ldots,f_{0,m})(x,y) \in \mathfrak{M}(r;k,m)$ and let*

$$\mathfrak{M}(r;k)^{l+1} \subset \mathfrak{M}(r;k)(\langle f_{0,i}, x\frac{\partial f_{0,i}}{\partial x_1}\rangle + \mathfrak{M}(r;k)\langle\frac{\partial f_{0,i}}{\partial y}\rangle) + \mathfrak{M}(r;k)^{l+2}$$

for $i = 1,\ldots,m$. Then f_0 is reticular $\mathcal{K}_{(m)}$-l-determined. Conversely if $f_0(x,y) \in \mathfrak{M}(r;k,m)$ is reticular $\mathcal{K}_{(m)}$-l-determined, then

$$\mathfrak{M}(r;k)^{l+1} \subset \langle f_{0,i}, x\frac{\partial f_{0,i}}{\partial x_1}\rangle_{\mathcal{E}(r;k)} + \mathfrak{M}(r;k)\langle\frac{\partial f_{0,i}}{\partial y}\rangle \quad \text{for } i = 1,\ldots,m.$$

We say that $f = (f_1,\ldots,f_m)(x,y,u)$, $g = (g_1,\ldots,g_m)(x,y,u) \in \mathcal{E}(r;k+n,m)$ are *reticular $(\mathcal{P}$-$\mathcal{K})_{(m)}$-equivalent* if there exist $\Phi_i \in \mathcal{B}_n(r;k+n)$ of the form:

$$\Phi_i(x,y,u) = (x\phi^i_1(x,y,t,u), \phi^i_2(x,y,t,u), \phi_3(u))$$

and a unit $\alpha = (\alpha_1,\ldots,\alpha_m)(x,y,u) \in \mathcal{E}(r;k+n,m)$ such that $g_i = \alpha_i \cdot f_i \circ \Phi_i$ for $i = 1,\ldots,m$. We write $\Phi = (\Phi_1,\ldots,\Phi_m)$ and call (Φ,α) a reticular $(\mathcal{P}$-$\mathcal{K})_{(m)}$-isomorphism from f to g.

We say that a map germ $f = (f_1,\ldots,f_m)(x,y,u) \in \mathfrak{M}(r;k+n,m)$ is *reticular $(\mathcal{P}$-$\mathcal{K})_{(m)}$-l-determined* if all map germ in $\mathfrak{M}(r;k+n,m)$ which has the same l-jet of f is reticular $(\mathcal{P}$-$\mathcal{K})_{(m)}$-equivalent to f.

Lemma 2.4 *Let* $f = (f_1, \dots, f_m)(x, y, u) \in \mathcal{E}(r; k+n, m)$ *and* $z = j^l f(0)$. *We write* $O^l_{r\mathcal{P}\text{-}\mathcal{K}_{(m)}}(z)$ *the submanifold of the image of the orbit of* f *under the reticular* t-$(\mathcal{P}$-$\mathcal{K})_{(m)}$ *equivalence relation. Then*

$$\begin{aligned} O^l_{r\mathcal{P}\text{-}\mathcal{K}_{(m)}}(z) = \pi_l((\langle f_1, x\frac{\partial f_1}{\partial x}\rangle_{\mathcal{E}(r;k+n)} + \mathfrak{M}(r; k+n)\langle \frac{\partial f_1}{\partial y}\rangle) \\ \times \cdots \times (\langle f_m, x\frac{\partial f_m}{\partial x}\rangle_{\mathcal{E}(r;k+n)} + \mathfrak{M}(r; k+n)\langle \frac{\partial f_m}{\partial y}\rangle) + \mathfrak{M}(n)\langle \frac{\partial f}{\partial u}\rangle). \end{aligned}$$

Lemma 2.5 (Cf., [10, Lemma 3.10]) *Let* $f = (f_1, \dots, f_m)(x, y, u) \in \mathfrak{M}(r; k+n, m)$ *and* l *be a non-negative integer. If*

$$\begin{aligned} \mathfrak{M}(r; k+n, m)^l \subset (\langle f_1, x\frac{\partial f_1}{\partial x}\rangle_{\mathcal{E}(r;k+n)} + \mathfrak{M}(r; k+n)\langle \frac{\partial f_1}{\partial y}\rangle) \times \cdots \times \\ (\langle f_m, x\frac{\partial f_m}{\partial x}\rangle_{\mathcal{E}(r;k+n)} + \mathfrak{M}(r; k+n)\langle \frac{\partial f_m}{\partial y}\rangle) + \mathfrak{M}(n)\langle \frac{\partial f}{\partial u}\rangle \\ + \mathfrak{M}(n)\mathfrak{M}(r; k+n, m)^l, \end{aligned} \tag{1}$$

then f *is reticular* $(\mathcal{P}$-$\mathcal{K})_{(m)}$-l*-determined.*

In convenience, we denote an unfolding of a function germ $f(x, y, u) \in \mathfrak{M}(r; k+n, m)$ by $F(x, y, t, u) \in \mathfrak{M}(r; k+m'+n, m)$.

Let $F(x, y, t, u) \in \mathfrak{M}(r; k+m'_1+n, m)$ and $G(x, y, s, u) \in \mathfrak{M}(r; k+m'_2+n, m)$ be unfoldings of $f(x, y, u) \in \mathfrak{M}(r; k+n, m)$.

A reticular t-$(\mathcal{P}$-$\mathcal{K})_{(m)}$-f*-morphism from* F *to* G is a pair (Φ, α), where $\Phi = (\Phi_1, \dots, \Phi_m)$ for $\Phi_i \in \mathfrak{M}(r; k+m'_2+n, r; k+m'_1+n)$ and $\alpha = (\alpha_1, \dots, \alpha_m)$ is a unit of $\mathcal{E}(r; k+m'_2+n, m)$ satisfying the following conditions:
(1) Φ_i can be written in the form:

$$\Phi_i(x, y, s, u) = (x\phi^i_1(x, y, s, u), \phi^i_2(x, y, s, u), \phi_3(s), \phi_4(s, u)),$$

(2) $\Phi_i|_{\mathbb{H}^r \times \mathbb{R}^{k+n}} = id_{\mathbb{H}^r \times \mathbb{R}^{k+n}}$, $\alpha_i|_{\mathbb{H}^r \times \mathbb{R}^{k+n}} \equiv 1$,
(3) $G_i(x, y, s, u) = \alpha_i(x, y, s, u) \cdot F_i \circ \Phi_i(x, y, s, u)$ for all $(x, y, s, u) \in (\mathbb{H}^r \times \mathbb{R}^{k+m'_2+n}, 0)$.

If there exists a reticular t-$(\mathcal{P}$-$\mathcal{K})_{(m)}$-f-morphism from F to G, we say that G *is reticular* t-$(\mathcal{P}$-$\mathcal{K})_{(m)}$-f*-induced from* F. If $m_1 = m_2$ and Φ is invertible, we call (Φ, α) *a reticular* t-$(\mathcal{P}$-$\mathcal{K})_{(m)}$-f*-isomorphism from* F *to* G and we say that F is reticular t-$(\mathcal{P}$-$\mathcal{K})_{(m)}$-f-equivalent to G.

We say that $F(x, y, t, u), G(x, y, t, u) \in \mathcal{E}(r; k+m'+n, m)$ are *reticular* t-$(\mathcal{P}$-$\mathcal{K})_{(m)}$*-equivalent* if there exist diffeomorphism germs

$$\Phi_i : (\mathbb{H}^r \times \mathbb{R}^{k+m'+n}, 0) \to (\mathbb{H}^r \times \mathbb{R}^{k+m'+n}, 0) \ \ (i = 1, \dots, m)$$

of the form

$$\Phi_i(x, y, t, q, z) = (x\phi^i_1(x, y, t, u), \phi^i_2(x, y, t, u), \phi_3(t), \phi_4(t, u))$$

and a unit $\alpha = (\alpha_1, \dots, \alpha_m) \in \mathcal{E}(r; k+m'+n, m)$ such that $G_i = \alpha_i \cdot F_i \circ \Phi_i$ for $i = 1, \dots, m$. We call (Φ, α) *a reticular* t-$(\mathcal{P}$-$\mathcal{K})_{(m)}$*-isomorphism from* F *to* G.

We say that function germs $F_0(x_1,\ldots,x_{r_1},y_1,\cdots,y_{k_1},t,q,z)\in\mathfrak{M}\,(r_1;k_1+1+n+1)$ and $F_1(x_1,\ldots,x_{r_2},y_1,\cdots,y_{k_2},t,q,z)\in\mathfrak{M}(r_2;k_2+1+n+1)$ are *stably reticular* t*-*$\mathcal{P}$*-*$\mathcal{K}$*-equivalent* if F_0 and F_1 are reticular t-$\mathcal{P}$-$\mathcal{K}$-equivalent after additions of linear forms of x of which all coefficients are not zero and non-degenerate quadratic forms in the variables y.

We define the notion of *stable reticular* t*-*$(\mathcal{P}$*-*$\mathcal{K})_{(m)}$*-equivalence* in the same way as the stable reticular t-$\mathcal{P}$-$\mathcal{K}$-equivalence.

Definition 2.6 We define stabilities of unfoldings. Let $F=(F_1,\ldots,F_m)(x,y,t,u)\in\mathfrak{M}(r;k+m'+n,m)$ be an unfolding of $f=(f_1,\ldots,f_m)(x,y,u)\in\mathfrak{M}(r;k+n,m)$.

We say that F is *reticular* t*-*$(\mathcal{P}$*-*$\mathcal{K})_{(m)}$*-stable* if the following condition holds: For any neighbourhood U of 0 in $\mathbb{H}^r\times\mathbb{R}^{k+m'+n}$ and any representative $\tilde{F}\in C^\infty(U,\mathbb{R}^m)$, there exists a neighbourhood $N_{\tilde{F}}$ of $\tilde{F}$ in C^∞-topology such that for any $\tilde{G}\in N_{\tilde{F}}$ there exist $(0,y^i,t^0,q^0,z^0)\in U$ for $i=1,\ldots,m$ such that $G=(G_1,\ldots,G_m)$ and F are reticular t-$(\mathcal{P}$-$\mathcal{K})_{(m)}$-equivalent where $G_i\in\mathfrak{M}(r;k+m'+n)$ is defined by $G_i(x,y,t,u)=\tilde{G}(x,y+y^i,t+t^0,u+u^0)-\tilde{G}(0,y,t^0,u^0)$.

We say that F is *a reticular* t*-*$(\mathcal{P}$*-*$\mathcal{K})_{(m)}$*-versal unfolding of* f if any unfolding of f is reticular t-$(\mathcal{P}$-$\mathcal{K})_{(m)}$-f-induced from F.

We say that F is *a reticular* t*-*$(\mathcal{P}$*-*$\mathcal{K})_{(m)}$*-universal unfolding of* f if m is minimal in reticular t-$(\mathcal{P}$-$\mathcal{K})_{(m)}$-versal unfoldings of f.

We say that F is *reticular* t*-*$(\mathcal{P}$*-*$\mathcal{K})_{(m)}$*-infinitesimally versal* if

$$\begin{aligned}\mathcal{E}(r;k+n,m)=\langle f_1,x\frac{\partial f_1}{\partial x},\frac{\partial f_1}{\partial y}\rangle_{\mathcal{E}(r;k+n)}\times\cdots\times\\ \langle f_m,x\frac{\partial f_m}{\partial x},\frac{\partial f_m}{\partial y}\rangle_{\mathcal{E}(r;k+n)}+\langle\frac{\partial f}{\partial u}\rangle_{\mathcal{E}(n)}+\langle\frac{\partial F}{\partial t}|_{t=0}\rangle_{\mathbb{R}}\end{aligned}$$

We say that F is *reticular* t*-*$(\mathcal{P}$*-*$\mathcal{K})_{(m)}$*-infinitesimally stable* if

$$\begin{aligned}\mathcal{E}(r;k+m'+n,m)=\langle F_1,x\frac{\partial F_1}{\partial x},\frac{\partial F_1}{\partial y}\rangle_{\mathcal{E}(r;k+m'+n)}\times\cdots\times\\ \langle F_m,x\frac{\partial F_m}{\partial x},\frac{\partial F_m}{\partial y}\rangle_{\mathcal{E}(r;k+m'+n)}+\langle\frac{\partial F}{\partial u}\rangle_{\mathcal{E}(m'+n)}+\langle\frac{\partial F}{\partial t}\rangle_{\mathcal{E}(m')}\end{aligned}$$

We say that F is *reticular* t*-*$(\mathcal{P}$*-*$\mathcal{K})_{(m)}$*-homotopically stable* if for any smooth path-germ $(\mathbb{R},0)\to\mathcal{E}(r;k+m'+n,m),\tau\mapsto F_\tau=(F_{1,\tau},\ldots,F_{m,\tau})$ with $F_0=F$, there exists a smooth path-germs $(\mathbb{R},0)\to\mathcal{B}(r;k+m'+n)\times\mathcal{E}(r;k+m'+n),\tau\mapsto(\Phi^i_\tau,\alpha^i_\tau)$ with $(\Phi^i_0,\alpha^i_0)=(id,1)$ and Φ^i_τ has the form

$$\Phi^i_\tau(x,y,t,u)=(x\phi^{i,1}_\tau(x,y,t,u),\phi^{i,2}_\tau(x,y,t,u),\phi^3_\tau(t),\phi^4_\tau(t,u))$$

such that each $(\Phi^i_\tau,\alpha^i_\tau)$ is a reticular t-$\mathcal{P}$-$\mathcal{K}$-isomorphism and $F_{i,\tau}=\alpha^i_\tau\cdot F_{i,0}\circ\Phi^i_\tau$ for $\tau\in(\mathbb{R},0)$ and $i=1,\ldots,m$.

Let U be a neighbourhood of 0 in $\mathbb{H}^r \times \mathbb{R}^{k+m'+n}$ and let $\tilde{F} = (\tilde{F}^1, \ldots, \tilde{F}^m) : U \to \mathbb{R}^m$ be a smooth map and l be a non-negative integer. We choose a neighbourhood U' of 0 in $\mathbb{R}^{km+m'+n}$ such that $(0, y^i, t, u) \in U$ for any $(y^1, \ldots, y^m, t, u) \in U$ and i. We define the smooth map germ

$$j^l_1 \tilde{F} : U' \longrightarrow J^l(r+k+n)$$

as the follow: For $(y^1, \ldots, y^m, t, u) \in U$ we set $j^l_1 \tilde{F}(y^1, \ldots, y^m, t, u)$ by the l-jet of the map germ $(\tilde{F}^1_{(y^1,t,u)}, \ldots, \tilde{F}^m_{(y^m,t,u)}) \in \mathfrak{M}(r; k+n, m)$ at 0, where $\tilde{F}^i_{(y^i,t,u)}$ is given by $\tilde{F}^i_{(y^i,t,u)}(x', y', u') = \tilde{F}(x', y^i + y', t, u + u') - \tilde{F}^i(0, y^i, t, u)$

Let $F(x, y, t, u) \in \mathfrak{M}(r; k+m'+n, m)$ be an unfolding of $f(x, y, u) \in \mathfrak{M}(r; k+n, m)$. Let l be a non-negative integer and $z = j^l f(0)$. We say that F is *reticular* t*-*$(\mathcal{P}\text{-}\mathcal{K})_{(m)}$*-*$l$*-transversal* if $j^l_1 \tilde{F}$ at 0 is transversal to $O^l_{r\mathcal{P}\text{-}\mathcal{K}_{(m)}}(z)$ for a representative $\tilde{F} \in C^\infty(U, \mathbb{R}^m)$ of F.

Lemma 2.7 (Cf., [10, Lemma 3.4]) *Let* $F(x, y, t, u) \in \mathfrak{M}(r; k+m'+n, m)$ *be an unfolding of* $f(x, y, u) \in \mathfrak{M}(r; k+n, m)$*. Then* F *is reticular* t*-*$(\mathcal{P}\text{-}\mathcal{K})_{(m)}$*-*$l$*-transversal unfolding of* f *if and only if*

$$\begin{aligned}\mathcal{E}(r; k+n, m) = \langle f_1, x\frac{\partial f_1}{\partial x}, \frac{\partial f_1}{\partial y}\rangle_{\mathcal{E}(r;k+n)} \times \cdots \times \\ \langle f_m, x\frac{\partial f_m}{\partial x}, \frac{\partial f_m}{\partial y}\rangle_{\mathcal{E}(r;k+n)} + \langle \frac{\partial f}{\partial u}\rangle_{\mathcal{E}(n)} + \langle \frac{\partial F}{\partial t}|_{t=0}\rangle_{\mathbb{R}} \\ +\mathfrak{M}(r; k+n)^{l+1}\mathcal{E}(r; k+n, m).\end{aligned}$$

Theorem 2.8 (Uniqueness of universal unfoldings)(Cf., [10, Theorem 3.13]) *Let* $F(x, y, u, t), G(x, y, u, t) \in \mathfrak{M}(r; k+n+m', m)$ *be unfoldings of* $f \in \mathfrak{M}(r; k+n, m)$*. If* F *and* G *are reticular* t*-*$(\mathcal{P}\text{-}\mathcal{K})_{(m)}$*-universal, then* F *and* G *are reticular* t*-*$\mathcal{P}$*-*$\mathcal{K}$*-*f*-equivalent.*

Theorem 2.9 (Cf., [10, Theorem 3.14]) *Let* $F(x, y, t, u) \in \mathfrak{M}(r; k+m'+n, m)$ *be an unfolding of* $f(x, y, u) \in \mathfrak{M}(r; k+n, m)$ *and let* f *is an unfolding of* $f_0(x, y) \in \mathfrak{M}(r; k, m)$*. Then following are equivalent.*

(1) *There exists a non-negative number* l *such that* f_0 *is reticular* $\mathcal{K}_{(m)}$*-*l*-determined and* F *is reticular* t*-*$(\mathcal{P}\text{-}\mathcal{K})_{(m)}$*-*$l'$*-transversal for* $l' \geq lm + l + m'$*,*
(2) F *is reticular* t*-*$(\mathcal{P}\text{-}\mathcal{K})_{(m)}$*-stable,*
(3) F *is reticular* t*-*$(\mathcal{P}\text{-}\mathcal{K})_{(m)}$*-versal,*
(4) F *is reticular* t*-*$(\mathcal{P}\text{-}\mathcal{K})_{(m)}$*-infinitesimally versal,*
(5) F *is reticular* t*-*$(\mathcal{P}\text{-}\mathcal{K})_{(m)}$*-infinitesimally stable,*
(6) F *is reticular* t*-*$(\mathcal{P}\text{-}\mathcal{K})_{(m)}$*-homotopically stable.*

3 Reticular Legendrian unfoldings

We review the theory of reticular Legendrian unfoldings. Let $J^1(\mathbb{R}^n, \mathbb{R})$ be the 1-jet bundle of functions in n-variables which may be considered as $\mathbb{R}^{2n+1}$ with a natural coordinate system $(q, z, p) = (q_1, \ldots, q_n, z, p_1, \ldots, p_n)$, where $(q_1, \ldots, q_n)$ is a coordinate system of $\mathbb{R}^n$. We equip the contact structure on $J^1(\mathbb{R}^n, \mathbb{R})$ defined by the canonical 1-form $\theta = dz - \sum_{i=1}^n p_i dq_i$. We have a natural projection $\pi : J^1(\mathbb{R}^n, \mathbb{R}) \to \mathbb{R}^n \times \mathbb{R}$ by $\pi(q, z, p) = (q, z)$.

We remark that: By the identification of the coordinate system $(u_1, \ldots, u_{n+1})$ and $(q_1, \ldots, q_n, z)$ on a manifold, the dimension of a manifold on which wavefronts propagate is higher 1 than n.

We also consider the 1-jet bundle $J^1(\mathbb{R} \times \mathbb{R}^n, \mathbb{R})$ and the canonical 1-form Θ on that space. Let $(t, q_1, \ldots, q_n)$ be the canonical coordinate system on $\mathbb{R} \times \mathbb{R}^n$ and

$$(t, q_1, \ldots, q_n, z, s, p_1, \ldots, p_n)$$

be the corresponding coordinate system on $J^1(\mathbb{R} \times \mathbb{R}^n, \mathbb{R})$. Then the canonical 1-form Θ is given by

$$\Theta = dz - \sum_{i=1}^{n} p_i dq_i - sdt = \theta - sdt.$$

We define the natural projection $\Pi : J^1(\mathbb{R}\times\mathbb{R}^n, \mathbb{R}) \to (\mathbb{R}\times\mathbb{R}^n)\times\mathbb{R}$ by $\Pi(t, q, z, s, p) = (t, q, z)$.

Let $\tilde{L}^{r,0}_{\sigma} = \{(t, q, z, s, p) \in J^1(\mathbb{R} \times \mathbb{R}^n, \mathbb{R}) | q_\sigma = p_{I_r - \sigma} = q_{r+1} = \cdots = q_n = s = z = 0, q_{I_r - \sigma} \geq 0\}$ for $\sigma \subset I_r = \{1, 2, \cdots, r\}$ and let $\mathbb{L}^r = \{(t, q, z, s, p) \in J^1(\mathbb{R} \times \mathbb{R}^n, \mathbb{R}) | q_1 p_1 = \cdots = q_r p_r = q_{r+1} = \cdots = q_n = s = z = 0, q_{I_r} \geq 0\}$ be a representative as a germ of the union of $\tilde{L}^{r,0}_{\sigma}$ for all $\sigma \subset I_r$.

We recall that $\tilde{L}^{r,0}_{\sigma}$ is a normalisation of the Legendrian submanifold consists of the particles generated by the *σ-corner* of hypersurface germ $V = \{(u_1, \ldots, u_n, 0) \in (\mathbb{R}^{n+1}, 0) | u_{I_r} \geq 0\}$ with an r-corner to conormal direction, where σ-corner is defined by $V \cap \{u_\sigma = 0\}$ and u is a local coordinate system of a manifold in which wavefronts propagate.

Definition 3.1 Let $C : (J^1(\mathbb{R} \times \mathbb{R}^n, \mathbb{R}), 0) \to (J^1(\mathbb{R} \times \mathbb{R}^n, \mathbb{R}), w)$ ($\Pi(w) = 0$) be a contact diffeomorphism germ. We say that C is a *$\mathcal{P}$-contact diffeomorphism germ* if C has the form:

$$C(t, q, z, s, p) = (t, q_t(q, z, p), z_t(q, z, p), h(t, q, z, s, p), p_t(q, z, p)). \tag{2}$$

We say a map germ $\mathcal{L} : (\mathbb{L}^r, 0) \to (J^1(\mathbb{R} \times \mathbb{R}^n, \mathbb{R}), w)$ $(\Pi(w) = 0)$ *a reticular Legendrian unfolding* if there exists a $\mathcal{P}$-contact diffeomorphism germ C such that $\mathcal{L}$ is the restriction of C to $\mathbb{L}^r$.

We call $\{\mathcal{L}(\tilde{L}^{r,0}_{\sigma})\}_{\sigma \subset I_r}$ *the unfolded contact regular r-cubic configuration of* $\mathcal{L}$.

In order to study perestroikas of wavefronts of reticular Legendrian unfoldings, we introduce the following equivalence relation. Let $\mathcal{L}_i : (\mathbb{L}^r, 0) \to (J^1(\mathbb{R} \times \mathbb{R}^n, \mathbb{R}), w_i), (i = 0, 1)$ be reticular Legendrian unfoldings. We say that $\mathcal{L}_1$ and $\mathcal{L}_2$ are *$\mathcal{P}$-Legendrian equivalent* if there exist a contact diffeomorphism germ

$$K : (J^1(\mathbb{R} \times \mathbb{R}^n, \mathbb{R}), w_1) \to (J^1(\mathbb{R} \times \mathbb{R}^n, \mathbb{R}), w_2)$$

of the form

$$\begin{aligned} K(t, q, z, s, p) = (\phi_1(t), \phi_2(t, q, z), \phi_3(t, q, z), \\ \phi_4(t, q, z, s, p), \phi_5(t, q, z, s, p)) \end{aligned} \tag{3}$$

and *a reticular r-diffeomorphism* Ψ on $(J^1(\mathbb{R} \times \mathbb{R}^n, \mathbb{R}), 0)$ such that $K \circ \mathcal{L}_1 = \mathcal{L}_2 \circ \Psi$, where a reticular r-diffeomorphism is defined by a contact diffeomorphism Ψ on $(J^1(\mathbb{R} \times \mathbb{R}^n, \mathbb{R}), 0)$ such that $t \circ \Psi$ depends only on t and Ψ preserves $\tilde{L}^{r,0}_{\sigma}$ for all $\sigma \subset I_r$.

We can construct generating families of reticular Legendrian unfoldings. A function germ $F(x,y,t,q,z) \in \mathfrak{M}(r;k+1+n+1)$ is called *$\mathcal{P}$-C-non-degenerate* if $\frac{\partial F}{\partial x}(0) = \frac{\partial F}{\partial y}(0) = 0$, $\frac{\partial F}{\partial z}(0) \neq 0$ and

$$x_1, \cdots, x_r, t, F, \frac{\partial F}{\partial x_1}, \cdots, \frac{\partial F}{\partial x_r}, \frac{\partial F}{\partial y_1}, \cdots, \frac{\partial F}{\partial y_k}$$

are independent on $(\mathbb{H}^k \times \mathbb{R}^{k+1+n+1}, 0)$.

Let $\mathcal{L}$ be a reticular Legendrian unfolding. A $\mathcal{P}$-C-non-degenerate function germ $F(x,y,t,q,z) \in \mathfrak{M}(r;k+1+n+1)$ is called *a generating family* of $\mathcal{L}$ if F is a generating family of the unfolded contact regular r-cubic configuration $\{\mathcal{L}(\tilde{L}^{r,0}_\sigma)\}_{\sigma \subset I_r}$, that is

$$\begin{aligned}\mathcal{L}(\tilde{L}^{r,0}_\sigma) = \{(t,q,z,\frac{\partial F}{\partial t}/(-\frac{\partial F}{\partial z}), \frac{\partial F}{\partial q}/(-\frac{\partial F}{\partial z})) \in (J^1(\mathbb{R}\times\mathbb{R}^n,\mathbb{R}), w) |\\ x_\sigma = F = \frac{\partial F}{\partial x_{I_r-\sigma}} = \frac{\partial F}{\partial y} = 0, x_{I_r-\sigma} \geq 0\}\end{aligned}$$

for all $\sigma \subset I_r$.

By [11, Theorem 0.8.5] we have that:

Theorem 3.2 (1) *For any reticular Legendrian unfolding $\mathcal{L} : (\mathbb{L}^r, 0) \to (J^1(\mathbb{R}\times\mathbb{R}^n,\mathbb{R}), w)$, there exists a function germ $F(x,y,t,q,z) \in \mathfrak{M}(r;k+1+n+1)$ which is a generating family of $\mathcal{L}$.*
(2) *For any $\mathcal{P}$-C-non-degenerate function germ $F(x,y,t,q,z) \in \mathfrak{M}(r;k+1+n+1)$ there exists a reticular Legendrian unfolding $\mathcal{L} : (\mathbb{L}^r, 0) \to (J^1(\mathbb{R}\times\mathbb{R}^n,\mathbb{R}), w)$ of which F is a generating family.*
(3) *Two reticular Legendrian unfolding are $\mathcal{P}$-Legendrian equivalent if and only if their generating families are stably reticular t-$\mathcal{P}$-$\mathcal{K}$-equivalent.*

4 Multi-reticular Legendrian unfolding

Let $\mathcal{L}_i : (\mathbb{L}^{r_i}, 0) \to (J^1(\mathbb{R}\times\mathbb{R}^n,\mathbb{R}), w_i) (i = 1, \dots, m)$ be reticular Legendrian unfoldings with $\Pi(w_i) = 0$ where $w_1, \dots, w_m$ are distinct. Then we call $\mathcal{L} = (\mathcal{L}_1, \dots, \mathcal{L}_m)$ *a multi-reticular Legendrian unfolding.*

Let $(\mathcal{L}_1, \dots, \mathcal{L}_m)$ and $(\mathcal{L}'_1, \dots, \mathcal{L}'_m)$ be multi-reticular Legendrian unfoldings. We say that they are reticular $\mathcal{P}_{(m)}$-Legendrian equivalent if there exist contact diffeomorphism germs

$$K_i : (J^1(\mathbb{R}\times\mathbb{R}^n,\mathbb{R}), w_i) \to (J^1(\mathbb{R}\times\mathbb{R}^n,\mathbb{R}), w'_i) \quad (i = 1, \dots, m)$$

of the form

$$\begin{aligned}&K_i(t,q,z,s,p) =\\ &\quad (\phi_1(t), \phi_2(t,q,z), \phi_3(t,q,z), \phi^i_4(t,q,z,s,p), \phi^i_5(t,q,z,s,p))\end{aligned}$$

and reticular $\mathcal{P}_{r_i}$-diffeomorphisms Ψ_i on $(\mathbb{L}^{r_i}, 0)$ such that $K_i \circ \mathcal{L}_i = \mathcal{L}'_i \circ \Psi_i$ for $i = 1, \dots, m$.

Let $(\mathcal{L}_1, \dots, \mathcal{L}_m)$ be a multi-reticular Legendrian unfoldings and $F_i \in \mathfrak{M}(r_i; k_i + 1 + n + 1)$ be generating families of $\mathcal{L}_i$. We call $F = (F_1, \dots, F_m)$ *a multi-generating family* of $(\mathcal{L}_1, \dots, \mathcal{L}_m)$.

We say that a map germs $(F_1,\dots,F_m)\in\mathfrak{M}(r;k+1+n+1)$ is $\mathcal{P}_{(m)}$-*C-non-degenerate* if all F_i are $\mathcal{P}$-C-non-degenerate and $(\frac{\partial F_1}{\partial t}/(-\frac{\partial F_1}{\partial z}),\frac{\partial F_1}{\partial q}/(-\frac{\partial F_1}{\partial z})),\dots,(\frac{\partial F_m}{\partial t}/(-\frac{\partial F_m}{\partial z}),\frac{\partial F_m}{\partial q}/(-\frac{\partial F_m}{\partial z}))$ are distinct.

By Theorem 0.8.5 in [11] we have that:

Proposition 4.1 (1) *For any multi-reticular Legendrian unfolding $\mathcal{L}$, there exists a multi-generating family of $\mathcal{L}$,*
(2) *For any $\mathcal{P}_{(m)}$-C-non-degenerate map germs $F=(F_1,\dots,F_m)$ $(F_i\in\mathfrak{M}(r_i;k_i+1+n+1))$, there exists a multi-reticular Legendrian unfolding of which F is a multi-generating family,*
(3) *Let $F=(F_1,\dots,F_m)$ and $F'=(F'_1,\dots,F'_m)$ be multi-generating families of multi-reticular Legendrian unfoldings $(\mathcal{L}_1,\dots,\mathcal{L}_m)$ and $(\mathcal{L}'_1,\dots,\mathcal{L}'_m)$ respectively. Then $(\mathcal{L}_1,\dots,\mathcal{L}_m)$ and $(\mathcal{L}'_1,\dots,\mathcal{L}'_m)$ are $\mathcal{P}_{(m)}$-Legendrian equivalent if and only if F and F' are stably reticular t-$(\mathcal{P}$-$\mathcal{K})_{(m)}$-equivalent.*

Definition 4.2 We define stabilities of multi-reticular Legendrian unfoldings. Let $\mathcal{L}=(\mathcal{L}_1,\dots,\mathcal{L}_m)$ be a multi-reticular Legendrian unfolding.
Stability: We say that $\mathcal{L}$ is *stable* if the following condition holds: Let $C^{0,i}\in C_T(J^1(\mathbb{R}\times\mathbb{R}^n,\mathbb{R}),0)$ $(i=1,\dots,m)$ be $\mathcal{P}$-contact embedding germs such that $C^{0,i}|_{\mathbb{L}^{r_i}}=\mathcal{L}_i$ and $\tilde{C}^{0,i}\in C_T(U_i,J^1(\mathbb{R}\times\mathbb{R}^n,\mathbb{R}))$ be a representative of $C^{0,i}$. Then there exist open neighbourhoods $N_{\tilde{C}^{0,i}}$ of $\tilde{C}^{0,i}$ in C^∞-topology for $i=1,\dots,m$ such that for any $\tilde{C}^i\in N_{\tilde{C}^{0,i}}$, there exist points $x_i=(T^i,0,\dots,0,P^i_{r+1},\dots,P^i_n)\in U_i$ such that the multi-reticular Legendrian unfolding $(\mathcal{L}'_{x_1},\dots,\mathcal{L}'_{x_m})$ and $\mathcal{L}$ are $\mathcal{P}_{(m)}$-Legendrian equivalent, where reticular Legendrian unfoldings $\mathcal{L}'_{x_i}$ are chosen that the reticular Legendrian unfolding $\tilde{C}^i|_{\mathbb{L}^{r_i}}:(\mathbb{L}^{r_i},x_i)\to(J^1(\mathbb{R}\times\mathbb{R}^n,\mathbb{R}),\tilde{C}^i(x_i))$ and $\mathcal{L}'_{x_i}$ are $\mathcal{P}$-Legendrian equivalent. We remark that the definition of stability is not depend on choices of $\mathcal{L}'_{x_i}$.
Homotopical stability: We say that $\mathcal{L}$ is *homotopically stable* if for any reticular $\mathcal{P}$-contact deformations $\bar{C}^i=\{C^i_\tau\}$ of $\mathcal{L}_i$, there exist one-parameter families of $\mathcal{P}$-Legendrian equivalences $\bar{K}^i=\{K^i_\tau\}$ on $(J^1(\mathbb{R}\times\mathbb{R}^n,\mathbb{R}),w_i)$ with $K^i_0=id$ of the form

$$K^i_\tau(t,q,z,s,p)=\\ (\phi^1_\tau(t),\phi^2_\tau(t,q,z),\phi^3_\tau(t,q,z),\phi^{4,i}_\tau(t,q,z,s,p),\phi^{5,i}_\tau(t,q,z,s,p)) \qquad (4)$$

and a one-parameter deformation of reticular $\mathcal{P}_{r_i}$-diffeomorphisms $\bar{\Psi}^i=\{\Psi^i_\tau\}$ such that $C^i_\tau=K^i_\tau\circ C^i_0\circ\Psi^i_\tau$ for t around 0 and $i=1,\dots,m$.
Infinitesimal stability: We say that $\mathcal{L}$ is *infinitesimally stable* if for any extension C^i of $\mathcal{L}_i$ and any infinitesimal $\mathcal{P}$-contact transformation v_i of C^i, there exist infinitesimal reticular $\mathcal{P}_{r_i}$-diffeomorphisms ξ_i and infinitesimal $\mathcal{P}$-Legendrian equivalences η_i at w_i of the form

$$\eta_i(t,q,z,s,p)=a_1(t)\frac{\partial}{\partial t}+a_2(t,q,z)\frac{\partial}{\partial q}+a_3(t,q,z)\frac{\partial}{\partial z}\\ +a^i_4(t,q,z,s,p)\frac{\partial}{\partial s}+a^i_5(t,q,z,s,p)\frac{\partial}{\partial p}$$

such that $v_i=C^i_*\xi_i+\eta_i\circ C^i$ for $i=1,\dots,m$.

We denote $C_T(U,J^1(\mathbb{R}\times\mathbb{R}^n,\mathbb{R}))^{(m)}=C_T(U,J^1(\mathbb{R}\times\mathbb{R}^n,\mathbb{R}))\times\cdots\times C_T(U,J^1(\mathbb{R}\times\mathbb{R}^n,\mathbb{R}))$ (m-products) and denote other notations analogously. We call an element $C=(C_1,\dots,C_m)\in C_T(J^1(\mathbb{R}\times\mathbb{R}^n,\mathbb{R}),0)^{(m)}$ *a $\mathcal{P}_{(m)}$-contact diffeomorphism germ* if $\Pi\circ C(0)=0$ and $C_1(0),\dots,C_m(0)$ are distinct.

Theorem 4.3 (Multi $\mathcal{P}$-Contact transversality theorem) *Let Q_i, $i = 1, 2, \ldots$ are submanifolds of $J^l_{C_T^\Theta}(U, J^1(\mathbb{R} \times \mathbb{R}^n, \mathbb{R}))^{(m)}$. Then the set*

$$T = \{C = (C_1, \ldots, C_m) \in C_T^\Theta(U, J^1(\mathbb{R} \times \mathbb{R}^n, \mathbb{R}))^{(m)} | \\ j^l C \textit{ is transversal to } Q_i \textit{ for all } i \in \mathbb{N}\}$$

is a residual set in $C_T^\Theta(U, J^1(\mathbb{R} \times \mathbb{R}^n, \mathbb{R}))^{(m)}$

This is directly proved by $\mathcal{P}$-Contact transversality theorem ([**?**, Theorem 0.4.5]).

We denote the ring $\mathcal{E}(1 + n + n, m)$ on the coordinates (t, q, p) by $\mathcal{E}_{t,q,p(m)}$ and denote other notations analogously.

Theorem 4.4 *Let $\mathcal{L} = (\mathcal{L}_1, \ldots, \mathcal{L}_m)$ be a multi-reticular Legendrian unfolding with a multi-generating family $F = (F_1, \ldots, F_m)$. Let $C = (C_1, \ldots, C_m) \in C_T^\Theta(J^1(\mathbb{R} \times \mathbb{R}^n, \mathbb{R}), 0)^{(m)}$ be an extension of $\mathcal{L}$. Then the following are equivalent.*
(u) *F is a reticular t-$(\mathcal{P}$-$\mathcal{K})_{(m)}$-stable unfolding of $F|_{t=0}$.*
(hs) *$\mathcal{L}$ is homotopically stable.*
(is) *$\mathcal{L}$ is infinitesimally stable.*
(a) *$\mathcal{E}_{t,q,p(m)} = B_0^{r_1} \times \cdots \times B_0^{r_m} + \langle 1, p_1 \circ C', \ldots, p_n \circ C' \rangle_{(\Pi \circ C')^* \mathcal{E}_{t,q,z}} + \langle s \circ C' \rangle_{\mathcal{E}_t}$, where $C' = C|_{z=s=0}$, $B_0^{r_i} = \langle q_1 p_1, \ldots, q_{r_i} p_{r_i}, q_{r_i+1}, \ldots, q_n \rangle_{\mathcal{E}_{t,q,p}}$, and $(\Pi \circ C')^* \mathcal{E}_{t,q,z}$ be the $\mathcal{E}_{t,q,z}$-submodule of $\mathcal{E}_{t,q,z(m)}$ such that $a.(f_1, \ldots, f_m) = (a(\Pi \circ C'_1(f_1)), \ldots, a(\Pi \circ C'_m(f_m))$ for $a \in \mathcal{E}_{t,q,z}$, $(f_1, \ldots, f_m) \in \mathcal{E}_{t,q,p(m)}$.*

We remark that sufficiently near multi-reticular Legendrian unfoldings of stable one are stable by the condition (a).

5 Genericity

In order to give a generic classification of multi-reticular Legendrian unfoldings, we reduce our investigation to a finitely dimensional jet space of $\mathcal{P}$-contact diffeomorphism germs.

Definition 5.1 Let $\mathcal{L} = (\mathcal{L}_1, \ldots, \mathcal{L}_m)$ be a multi-reticular Legendrian unfolding. We say that $\mathcal{L}$ is l-determined if the following condition holds: For any extension $C_i \in C_T(J^1(\mathbb{R} \times \mathbb{R}^n, \mathbb{R}), 0)$ of $\mathcal{L}_i$, the multi-reticular Legendrian unfolding $(C'_1|_{\mathbb{L}^{r_1}}, \ldots, C'_m|_{\mathbb{L}^{r_m}})$ and $\mathcal{L}$ are $\mathcal{P}_{(m)}$-Legendrian equivalent for all $C'_i \in C_T(J^1(\mathbb{R} \times \mathbb{R}^n, \mathbb{R}), 0)$ satisfying that $j^l C_i(0) = j^l C'_i(0)$ for $i = 1, \ldots, m$.

As Lemma **??**, we may consider the following other definition of finitely determinacy of reticular Legendrian maps:
(1) The definition given by replacing $C_T(J^1(\mathbb{R} \times \mathbb{R}^n, \mathbb{R}), 0)$ to $C_T^\Theta(J^1(\mathbb{R} \times \mathbb{R}^n, \mathbb{R}), 0)$.
(2) The definition given by replacing $C_T(J^1(\mathbb{R} \times \mathbb{R}^n, \mathbb{R}), 0)$ to $C_T^Z(J^1(\mathbb{R} \times \mathbb{R}^n, \mathbb{R}), 0)$.
(3) The definition given by replacing $C_T(J^1(\mathbb{R} \times \mathbb{R}^n, \mathbb{R}), 0)$ to $C_T^{\Theta,Z}(J^1(\mathbb{R} \times \mathbb{R}^n, \mathbb{R}), 0)$.
Then the following holds by Proposition 0.5.2 in [**?**]:

Proposition 5.2 *Let $\mathcal{L}$ be a multi-reticular Legendrian unfolding. Then*
(A) *If $\mathcal{L}$ is l-determined of the original definition, then $\mathcal{L}$ is l-determined of the definition* (1).

(B) *If $\mathcal{L}$ is l-determined of the definition* (1)*, then $\mathcal{L}$ is l-determined of the definition* (3)*.*
(C) *If $\mathcal{L}$ is $(l+1)$-determined of the definition* (3)*, then $\mathcal{L}$ is l-determined of the definition* (2)*.*
(D) *If $\mathcal{L}$ is l-determined of the definition* (2)*, then $\mathcal{L}$ is l-determined of the original definition.*

Theorem 5.3 *Let $\mathcal{L} = (\mathcal{L}_1, \dots, \mathcal{L}_m)$ be a multi reticular Legendrian unfolding. If $\mathcal{L}$ is infinitesimally stable then $\mathcal{L}$ is $(n+5)$-determined.*

Let $J^l(2n+3, 2n+3)^{(m)}$ be the set of multi-l-jets of map germs from $(J^1(\mathbb{R}\times\mathbb{R}^n, \mathbb{R}), 0)$ to $J^1(\mathbb{R}\times\mathbb{R}^n, \mathbb{R})$ and $tC^l(n,m)$ be the Lie group in $J^l(2n+3, 2n+3)^{(m)}$ consists of multi-l-jets of $\mathcal{P}$-contact embedding germs. Let $L^l(2n+3)^{(m)}$ be the Lie group consists of multi-l-jets of diffeomorphism germs on $(J^1(\mathbb{R}\times\mathbb{R}^n, \mathbb{R}), 0)$.

We consider the Lie subgroup $rtLe^l(n,m)$ of $L^l(2n+3)^{(m)} \times L^l(2n+3)$ consists of multi-l-jets of reticular $\mathcal{P}_{r_i}$-diffeomorphisms on the source space and l-jets of $\mathcal{P}$-Legendrian equivalences of Π at 0:

$$\begin{aligned}&rtLe^l(n,m)\\ &\quad= \{(j^l\Psi_1(0), \dots, j^l\Psi_m(0), j^lK(0)) \in L^l(2n+3)^{(m)} \times L^l(2n+3) \mid\\ &\quad\Psi_i \text{ is a reticular } \mathcal{P}_{r_i}\text{-diffeomorphism on } (J^1(\mathbb{R}\times\mathbb{R}^n, \mathbb{R}), 0),\\ &\quad K \text{ is a } \mathcal{P}\text{-Legendrian equivalence of } \Pi\}.\end{aligned}$$

The group $rtLe^l(n,m)$ acts on $J^l(2n+3, 2n+3)$ and $tC^l(n,m)$ is invariant under this action. Let $C = (C_1, \dots, C_m)$ be a $\mathcal{P}_{(m)}$-contact diffeomorphism germ from $(J^1(\mathbb{R}\times\mathbb{R}^n, \mathbb{R}), 0)$ to $J^1(\mathbb{R}\times\mathbb{R}^n, \mathbb{R})$ and set $z_i = j^lC_i(0)$, $\mathcal{L}_i = C_i|_{\mathbb{L}^{r_i}}$, $\mathcal{L} = (\mathcal{L}_1, \dots, \mathcal{L}_m)$, $z = (z_1, \dots, z_m,)$. We denote the orbit $rtLe^l(n,m)\cdot z$ by $[z]$. Then

$$\begin{aligned}[z] = \{j^lC'(0) \in tC^l(n,m) \mid \mathcal{L} \text{ and } &(C'_1|_{\mathbb{L}^{r_1}}, \dots, C'_m|_{\mathbb{L}^{r_m}})\\ &\text{are } \mathcal{P}_{(m)}\text{-Legendrian equivalent}\}.\end{aligned}$$

We denote by VI_C the vector space consists of infinitesimal $\mathcal{P}_{(m)}$-contact transformation germs of C and denote by VI^0_C the subspace of VI_C consists of germs which vanish on 0. We denote by $VL_{J^1(\mathbb{R}\times\mathbb{R}^n,\mathbb{R})}$ by the vector space consists of infinitesimal $\mathcal{P}$-Legendrian equivalences on Π and denote by $VL^0_{J^1(\mathbb{R}\times\mathbb{R}^n,\mathbb{R})}$ by the subspace of $VL_{J^1(\mathbb{R}\times\mathbb{R}^n,\mathbb{R})}$ consists of germs which vanish at 0.

We denote by $V^0_{\mathbb{L}^r}$ the vector space consists of infinitesimal reticular r-diffeomorphisms on $(J^1(\mathbb{R}\times\mathbb{R}^n, \mathbb{R}), 0)$ which vanishes at 0 and set $V^0_{\mathbb{L}} = V^0_{\mathbb{L}^{r_1}} \times V^0_{\mathbb{L}^{r_m}}$. We have that by Lemma **??**:

$$\begin{aligned}VI^0_C = \{(v_1, \dots, v_m) | v_i : (J^1(\mathbb{R}\times\mathbb{R}^n, \mathbb{R}), 0) &\to T(J^1(\mathbb{R}\times\mathbb{R}^n, \mathbb{R})),\\ v_i = X_{f_i} \circ C_i &\text{ for some } f \in \mathfrak{M}^2_{t,q,z,p}\},\end{aligned}$$

$$\begin{aligned}VL^0_{J^1(\mathbb{R}\times\mathbb{R}^n,\mathbb{R})} = \{\eta \in X(J^1(\mathbb{R}\times\mathbb{R}^n, \mathbb{R}), 0) \mid &\eta = X_H \text{ for some}\\ \mathcal{P}\text{-fiver preserving function germ } &H \in \mathfrak{M}^2_{J^1(\mathbb{R}\times\mathbb{R}^n,\mathbb{R})}\},\end{aligned}$$

$$\begin{aligned}V^0_{\mathbb{L}} = \{(\xi_1, \dots, \xi_m) | \xi_i \in X(J^1(\mathbb{R}\times\mathbb{R}^n, \mathbb{R}), 0),\ &\xi_i = X_{g_i} \text{ for some}\\ &g_i \in B^{r_i} \cap \mathfrak{M}^2_{J^1(\mathbb{R}\times\mathbb{R}^n,\mathbb{R})}\}.\end{aligned}$$

We define the homomorphism $tC : \mathfrak{M}_{J^1(\mathbb{R}\times\mathbb{R}^n,\mathbb{R})}VI_{\mathbb{L}} \to VI^0_C$ by $tC(v) = (C_{1*}v_1, \ldots, C_{m*}v_m)$ and define the homomorphism $wC : VL^0_{J^1(\mathbb{R}\times\mathbb{R}^n,\mathbb{R})} \to VI^0_C$ by $wC(\eta) = (\eta \circ C_1, \ldots, \eta \circ C_m)$.

We denote VI^l_C the subspace of VI_C consists of infinitesimal $\mathcal{P}$-contact transformation germs of C whose l-jets are 0:

$$VI^l_C = \{(v_1, \ldots, v_m) \in VI_C \mid j^l v_i(0) = 0\}.$$

For $\tilde{C} = (\tilde{C}_1, \ldots, \tilde{C}_m) \in C_T(U, J^1(\mathbb{R}\times\mathbb{R}^n,\mathbb{R}))^{(m)}$, we define the continuous map $j^l_0\tilde{C} : U \to tC^l(n,m)$ by x to the l-jet of $(\tilde{C}_{1x}, \ldots, \tilde{C}_{mx})$.

Theorem 5.4 *Let $\mathcal{L} = (\mathcal{L}_1, \ldots, \mathcal{L}_m)$ be a reticular Legendrian unfolding. Let C_i be an extension of $\mathcal{L}_i$ and $l \geq (n+2)^2$. We set $C = (C_1, \ldots, C_m)$. Then the followings are equivalent:*
(s) *$\mathcal{L}$ is stable.*
(t) *$j^l_0 C$ is transversal to $[j^l_0 C(0)]$.*
(a') $\mathcal{E}_{t,q,p(m)} = B_0^{r_1} \times \cdots \times B_0^{r_m} + \langle 1, p_1 \circ C', \ldots, p_n \circ C'\rangle_{(\Pi\circ C')^*\mathcal{E}_{t,q,z}} + \langle s \circ C'\rangle_{\mathcal{E}_t} + \mathfrak{M}^l_{t,q,p(m)}$, *where $C' = C|_{z=s=0}$ and $B_0^{r_i} = \langle q_1p_1, \ldots, q_{r_i}p_{r_i}, q_{r_i+1}, \ldots, q_n\rangle_{\mathcal{E}_{t,q,p}}$,*
(a) $\mathcal{E}_{t,q,p(m)} = B_0^{r_1} \times \cdots \times B_0^{r_m} + \langle 1, p_1 \circ C', \ldots, p_n \circ C'\rangle_{(\Pi\circ C')^*\mathcal{E}_{t,q,z}} + \langle s \circ C'\rangle_{\mathcal{E}_t}$,
(is) *$\mathcal{L}$ is infinitesimally stable,*
(hs) *$\mathcal{L}$ is homotopically stable,*
(u) *A multi-generating family F of $\mathcal{L}$ is reticular t-$(\mathcal{P}$-$\mathcal{K})_{(m)}$-stable unfolding of $F|_{t=0}$.*

Let $\mathcal{L} = (\mathcal{L}_1, \ldots, \mathcal{L}_m)$ be a stable multi-reticular Legendrian unfolding. We say that $\mathcal{L}$ is *simple* if there exists a representative $\tilde{C} \in C_T(U, J^1(\mathbb{R}\times\mathbb{R}^n,\mathbb{R}))^{(m)}$ of a extension of $\mathcal{L}$ such that $\{\tilde{C}_x | x \in U\}$ is covered by finite orbits $[C_1], \ldots, [C_m]$ for some multi-$\mathcal{P}$-contact diffeomorphism germs $C_1, \ldots, C_m \in C_T(U, J^1(\mathbb{R}\times\mathbb{R}^n,\mathbb{R}))^{(m)}$.

Proposition 5.5 *Let $\mathcal{L} = (\mathcal{L}_1, \ldots, \mathcal{L}_m)$ be a stable multi-reticular Legendrian unfolding. Then $\mathcal{L}$ is simple if and only if all reticular Legendrian unfoldings $\mathcal{L}_i$ are simple for $i = 1, \ldots, m$.*

In order to classify generic multi-reticular Legendrian unfoldings, we classify stable multi-unfoldings $F = (F_1, \ldots, F_m)$ $(n \leq 2, m \geq 2)$ of $f = (f_1, \ldots, f_m)$ and $f_0 = (f_{0,1}, \ldots, f_{0,m})$ satisfying the condition: the reticular $(\mathcal{P}$-$\mathcal{K})_{(m-1)}$-codimension of $(f_1, \ldots, \check{f}_i, \ldots, , f_m) = 0$ for any i.

Let a stable multi-unfoldings $F = (F_1, \ldots, F_m)$ $(n \leq 2, m \geq 2)$ of $f = (f_1, \ldots, f_m)$ and $f_0 = (f_{0,1}, \ldots, f_{0,m})$ satisfying the condition be given. Then each F_i is a reticular t-$\mathcal{P}$-$\mathcal{K}$-stable unfolding of f_i, there exist monomials $\varphi_{i,1}, \ldots, \varphi_{i,\mu_i} \in \mathfrak{M}(r_i;k_i)$ such that they consist a basis of $Q_{f_{0,i}} = \langle f_{0,i}, x\frac{\partial f_{0,i}}{\partial x}, \frac{\partial f_{0,i}}{\partial y}\rangle_{\mathcal{E}(r_i;k_i)}$ and $\varphi_{i,1}$ has the maximal degree, and $\varphi_{i,\mu_i} = 1$. Then we have that $\mu_1 + \cdots \mu_m \leq n+2$. Since $\sum_{i=1}^m \mu_i \leq n+2 \leq 4$, we have that all $f_{0,i}$ are simple singularities. Therefore f_0 is stably reticular $\mathcal{K}_{(m)}$-equivalent to one of the multi-germs in the following list:
$n = 1$;
$m = 1$; y^2, y^3, y^4, x^2, x^3, $\pm xy + y^3$,
$m = 2$; (y^2, y^2), (y^2, y^3), (y^2, x^2),
$m = 3$; (y^2, y^2, y^2),
$n = 2$;

$m=1;\ y^2,\ y^3,\ y^4,\ y^5,\ y_1^2 \pm y_2^2,\ x^2,\ x^3,\ x^4,\ \pm xy+y^3,\ xy+y^4,\ x^2+y^3,$
$m=2; (y^2,y^2),\ (y^2,y^3),\ (y^2,y^4),\ (y^3,y^3),\ (y^2,x^2),\ (y^2,x^3),\ (y^2,\pm xy+y^3),\ (x^2,x^2),$
$m=3;\ (y^2,y^2,y^2),\ (y^2,y^2,y^3),\ (y^2,y^2,x^2),$
$m=4;\ (y^2,y^2,y^2,y^2).$

We construct a reticular $(\mathcal{P}\text{-}\mathcal{K})_{(m)}$-versal unfolding for each multi-germ by the usual method. Then the corresponding list is as follows:
$n=1;$
(1) $(y^2+u_{1,1}, y^2+u_{2,1})$
(2) $(y^2+u_{1,1}, y^3+u_{2,1}y+u_{2,2})$
(3) $(y^2+u_{1,1}, x^2+u_{2,1}x+u_{2,2})$
(4) $(y^2+u_{1,1}, y^2+u_{2,1}, y^2+u_{3,1})$
$n=2;$
(6) $(y^2+u_{1,1}, y^2+u_{2,1})$
(7) $(y^2+u_{1,1}, y^3+u_{2,1}y+u_{2,2})$
(8) $(y^2+u_{1,1}, y^4+u_{2,1}y^2+u_{2,2}y+u_{2,3})$
(9) $(y^3+u_{1,1}y+u_{1,2}, y^3+u_{2,1}y+u_{2,2})$
(10) $(y^2+u_{1,1}, x^2+u_{2,1}x+u_{2,2})$
(11) $(y^2+u_{1,1}, x^3+u_{2,1}x^2+u_{2,2}x+u_{2,3})$
(12) $(y^2+u_{1,1}, \pm xy+y^3+u_{2,1}y^2+u_{2,2}y+u_{2,3})$
(13) $(x^2+u_{1,1}x+u_{1,2}, x^2+u_{2,1}x+u_{2,2})$
(14) $(y^2+u_{1,1}, y^2+u_{2,1}, y^2+u_{3,1})$
(15) $(y^2+u_{1,1}, y^2+u_{2,1}, y^3+u_{3,1}y+u_{3,2})$
(16) $(y^2+u_{1,1}, y^2+u_{2,1}, x^2+u_{3,1}x+u_{3,2})$
(17) $(y^2+u_{1,1}, y^2+u_{2,1}, y^2+u_{3,1}, y^2+u_{4,1})$.

In the case that the reticular $(\mathcal{P}\text{-}\mathcal{K})_{(m)}$-codimension of $f=0$, the F is stably reticular t-$(\mathcal{P}\text{-}\mathcal{K})_{(m)}$-equivalent to $G=(G_1,\dots,G_m)$, where $G_i(x,y,t,q,z)=f_{0,i}(x,y)+u_{i,1}\varphi_1(x,y)+\cdots+u_{i,\mu_i}\varphi_{i,\mu_i}(x,y)-z$ for $i=1,\dots,m-1$, $G_m(x,y,t,q,z)=f_{0,i}(x,y)+u_{m,1}\varphi_1(x,y)+\cdots+u_{m,\mu_{m-1}}\varphi_{m,\mu_{m-1}}(x,y)-z$, and $(q_1,\dots,q_n,z)=(u_{1,1},\dots,u_{1,\mu_1},\dots,u_{m,1},\dots,u_{m,\mu_{m-1}},u_1,\dots,u_\mu,z)$.

In the case that the reticular $(\mathcal{P}\text{-}\mathcal{K})_{(m)}$-codimension of $f=1$, F is stably reticular t-$(\mathcal{P}\text{-}\mathcal{K})_{(m)}$-equivalent to $(F'_1,\dots,F'_m)$, where
(1) $F'_1=f_{0,1}(x,y)+(t+a(q,z))\varphi_1(x,y)+u_{1,1}\varphi_2(x,y)\cdots+u_{1,\mu_1-1}\varphi_{1,\mu_1}(x,\,y)-z,$
(2) $F'_i=f_{0,i}(x,y)+u_{i,1}\varphi_1(x,y)+\cdots+u_{i,\mu_i}\varphi_{i,\mu_i}(x,y)-z$ for $1<i<m$,
(3) $F'_m=f_{0,m}(x,y)+u_{m,1}\varphi_{m,1}(x,y)+\cdots+u_{m,\mu_m-1}\varphi_{m,\mu_m-1}(x,y)-z,$
(4) $(q_1,\dots,q_n,z)=(u_{1,1},\dots,u_{1,\mu_1-1},u_{2,1},\dots,u_{2,\mu_2},\dots,u_{m,1},\dots,u_{m,\mu_{m-1}},\ u_1,\dots,u_\mu,z),$
(5) $\frac{\partial a}{\partial u_{1,j}}(0)=0$ for $j=1,\dots,\mu_1-1$, $\frac{\partial a}{\partial z}(0)=0$, and $(\frac{\partial^2 a}{\partial u_i\partial u_j}(0))_{i,j=1,\dots,\mu}$ is non-degenerate.

We denote the linear part of a by $v=v_2+\cdots+v_m$, where v_i depends only on $u_{i,1},\dots,u_{i,\mu_i}$. Then we may reduce a to

$$a=t\pm u_{2,1}\pm\cdots\pm u_{m,1}\pm u_1^2\pm\cdots u_\mu^2.$$

Then F is stably reticular t-$(\mathcal{P}\text{-}\mathcal{K})_{(m)}$-equivalent to one of the following list:
$n=1;$

$m=2;$

${}^0({}^0A_1{}^0A_1):\quad (y^2+q-z, y^2-z);$
${}^1({}^0A_1{}^0A_1):\quad (y^2+t\pm q^2-z, y^2-z);$

${}^1({}^0A_1{}^0A_2):\ \ (y^2+t\pm q-z, y^3+qy-z);$
${}^1({}^0A_1{}^0B_2):\ \ (y^2+t\pm q-z, x^2+qx-z).$

$m=3;$

${}^1({}^0A_1{}^0A_1{}^0A_1):\ \ (y^2+t-z, y^2+q-z, y^2-q-z).$

$n=2;$

$m=2;$

${}^1({}^0A_1{}^0A_1):\ \ (y^2+t\pm q_1^2\pm q_2^2-z, y^2-z);$
${}^0({}^0A_1{}^0A_2):\ \ (y^2+t\pm q_1-z, y^3+q_1y-q_2-z);$
${}^1({}^0A_1{}^0A_2):\ \ (y^2+t\pm q_1\pm q_2^2-z, y^3+q_1y-z);$
${}^0({}^0A_1{}^0B_2):\ \ (y^2+q_1-z, x^2+q_2x-z);$
${}^1({}^0A_1{}^0A_3):\ \ (y^2+t\pm q_1-z, y^4+q_1y^2+q_2y-z);$
${}^1({}^0A_2{}^0A_2):\ \ (y^3+(t\pm q_1)y+q_2-z, y^3+q_1y-z);$
${}^1({}^0A_1{}^0B_2):\ \ (y^2+t\pm q_1\pm q_2^2-z, x^2+q_1x-z);$
${}^1({}^0A_1B_3):\ \ (y^2+t\pm q_1-z, x^3+q_1x^2+q_2x-z);$
${}^1({}^0A_1{}^0C_3^{\pm}):\ \ (y^2+t+q_1-z, \pm xy+y^3+q_1y^2+q_2y-z);$
${}^1({}^0B_2{}^0B_2):\ \ (x^2+(t\pm q_1)x+q_2-z, x^2+q_1x-z).$

$m=3;$

${}^0({}^0A_1{}^0A_1{}^0A_1):\ \ (y^2+q_1-z, y^2+q_2-z, y^2-z);$
${}^1({}^0A_1{}^0A_1{}^0A_1):\ \ (y^2+t\pm q_1\pm q_2^2-z, y^2+q_1-z, y^2-z);$
${}^1({}^0A_1{}^0A_1{}^0A_2):\ \ (y^2+t\pm q_1-z, y^2-z, y^3+q_1y+q_2-z);$
${}^1({}^0A_1{}^0A_1{}^0B_2):\ \ (y^2+t\pm q_1-z, y^2-z, x^2+q_1x+q_2-z).$

$m=4;$

${}^1({}^0A_1{}^0A_1{}^0A_1{}^0A_1):\ \ (y^2+t\pm q_1\pm q_2-z, y^2+q_1-z, y^2+q_2-z, y^2-z).$

Theorem 5.6 *Let $r_i=0$ or 1 for $i=1,\dots,m$ and $n\leq 2$. Let U be a neighborhood of 0 in $J^1(\mathbb{R}\times\mathbb{R}^n,\mathbb{R})$. Then there exists a residual set $O\subset C_T^{\Theta}(U,J^1(\mathbb{R}\times\mathbb{R}^n,\mathbb{R}))^{(m)}$ such that for any $\tilde{C}=(\tilde{C}_1,\dots,\tilde{C}_m)\in O$ and $x=(x_1,\dots,x_m)\in U^{(m)}$, the multi-reticular Legendrian unfolding $(\tilde{C}_{1x_1}|_{\mathbb{L}^{r_1}},\dots,\tilde{C}_{mx_m}|_{\mathbb{L}^{r_m}})$ is stable and have a multi-generating family which is stably reticular t-$(\mathcal{P}$-$\mathcal{K})_{(m)}$-equivalent for one of the types in the above list.*

Proof. Let $X_i=(X_{i,1},\cdots,X_{i,m})$ $(i=1,\dots,s)$ be all simple singularities with $\sum_{j=1}^m r$-$\mathcal{K}$-codim $X_{i,j}\leq n+2$, that is each $X_{i,j}$ is one of simple singularities A,B,C. For $i=1,\dots,s$ let F_{X_i} be $\mathcal{P}_{(m)}$-C-non-degenerate map germ which is unfolding of X_i. We choose an extension ${C^0}_{X_i}$ and ${C^1}_{X_i}$ of multi-reticular Legendrian unfoldings with multi-generating families ${F^0}_{X_i}$ and ${F^1}_{X_i}$ respectively. Let $l>16$. We define that

$$O'=\{\tilde{C}\in C_T^{\Theta}(U,J^1(\mathbb{R}\times\mathbb{R}^n,\mathbb{R}))^{(m)}\ |j_0^l\tilde{C} \text{ is transversal to } [(j^l{C^j}_{X_i}(0)] \text{ for all } i=1,\dots,s \text{ and } j=0,1\}.$$

Let $X_i'=(X_{i,1},\dots,X_{i,m-1})$ $(i=1,\dots,s')$ be all simple singularities with $\sum_{j=1}^{m-1} r$-$\mathcal{K}$-codim $X_{i,j}\leq n+1$. We choose ${C^0}_{X_i'}\in C_T^{\Theta}(U,J^1(\mathbb{R}\times\mathbb{R}^n,\mathbb{R}))^{(m-1)}$ analogous way. Then we define that

$$O''=\{\tilde{C}\in C_T^{\Theta}(U,J^1(\mathbb{R}\times\mathbb{R}^n,\mathbb{R}))^{(m)}\ |(j_0^l\tilde{C}_1,\dots,\check{j_0^l\tilde{C}_j},\dots,j_0^l\tilde{C}_m) \text{ is transversal to } [(j^l{C^0}_{X_i'}(0)] \text{ for all } i=1\cdots,s'\}.$$

Then O' and O'' are residual sets. We set

$$Y = \{j^l C(0) \in tC^l(n,m) \mid \text{the codimension of } [j^l C(0)] > 2n+4\}.$$

Then Y is an algebraic set in $tC^l(n,m)$. Therefore we can define that

$$O''' = \{\tilde{C} \in C_T^{\Theta}(U, J^1(\mathbb{R} \times \mathbb{R}^n, \mathbb{R}))^{(m)} \mid j_0^l \tilde{C} \text{ is transversal to } Y\}.$$

Then Y has codimension $> 2n+4$ because all $\mathcal{P}_{(m)}$-contact diffeomorphism germ with $j^l C(0) \in Y$ adjoin to the above list which are simple. Therefore

$$O''' = \{\tilde{C} \in C_T^{\Theta}(U, J^1(\mathbb{R} \times \mathbb{R}^n, \mathbb{R}))^{(m)} \mid j_0^l \tilde{C}(U^{(m)}) \cap Y = \emptyset\}.$$

Then the set $O = O' \cap O'' \cap O'''$ has the required condition. q.e.d.



Figure 2: ${}^0({}^0A_1{}^0A_1)$



Figure 3: ${}^0({}^0A_1{}^0A_1{}^0A_1)$



Figure 4: ${}^0({}^0A_1{}^0B_2)$



Figure 5: ${}^0({}^0A_1{}^0A_2)$

References

[1] V.I.Arnold,S.M. Gusein-Zade & A.N. Varchenko, *Singularities of differential maps,* Vol.I, Birkhäuser, Basel,1986.



Figure 6: $^1(^0A_1{}^0A_1)$



Figure 7: $^1(^0A_1{}^0A_2)$



Figure 8: $^1(^0A_2{}^0A_2)$



Figure 9: $^1(^0A_1{}^0B_2)$



Figure 10: $^1(^0A_1{}^0A_3)$



Figure 11: $^{1}(^{0}A_{1}{}^{0}B_{3})$



Figure 12: $^{1}(^{0}A_{1}{}^{0}C_{3}^{+})$



Figure 13: $^{1}(^{0}A_{1}{}^{0}C_{3}^{-})$



Figure 14: $^{1}(^{0}B_{2}{}^{0}B_{2})$



Figure 15: $^1(^0A_1{}^0A_1{}^0A_1)$



Figure 16: $^1(^0A_1{}^0A_1{}^0A_2)$



Figure 17: $^1(^0A_1{}^0A_1{}^0B_2)$



Figure 18: $^1(^0A_1{}^0A_1{}^0A_1{}^0A_1)$

[2] S.Izumiya and G.T.Kossioris, *Semi-local Classification of Geometric Singularities for Hamilton-Jacobi Equations*, J. of Differential Equations., **118**(1995), 166-193

[3] S.Izumiya, *Perestroikas of Optical Wave fronts and Graphlike Legendrian Unfoldings* , J. Differential Geom., **38**(1993), 485-500

[4] Th:bröcker, *Differentiable Germs and Catastrophes*, London Math. Soc. Lecture Note ser., 17 Cambridge Univ. Press, 1975.

[5] K.Jänich, *Caustics and catastrophes*, Math. Ann., 209(1974), p.161-180.

[6] T.Tsukada, *Reticular Lagrangian, Legendrian Singularities and their applications*, PhD Thesis, Hokkaido University, 1999.

[7] T.Tsukada, *Reticular Lagrangian Singularities*, The Asian J. of Math., vol 1, **3**(1997), 572-622.

[8] T.Tsukada, *Reticular Legendrian Singularities*, The Asian J. of Math., **1**(2001), 109-127.

[9] T.Tsukada, *Genericity of Caustics and Wavefronts on an r-corner*, in preparation.

[10] T.Tsukada, *A generic classification of function germs with respect to the reticular t-$\mathcal{P}$-$\mathcal{K}$-equivalence*, Hokkaido Math. J., 38, 177-203, 2009.

[11] T.Tsukada, *Bifurcations of Wavefronts on an r-corner*, in preparation.

[12] G.Wassermann, *Stability of unfolding*, Lecture note in mathematics, **393**.

[13] G.Wassermann, *Stability of unfolding in space and time*, Acta Math., **135**(1975), 57-128.

[14] V.M.Zakalyukin, *Reconstruction of fronts and caustics depending on a parameter and versality of mappings*, J. of Soviet Math., **27**(1984), 2713-2735